

Patient Instructions For Endovenous Laser Ablation

Please avoid aspirin products for 7 days prior to having the procedure. If you are on blood thinning medication (i.e. Plavix or Coumadin) please notify us (the risk is increased bruising). If pain medication is needed before the procedure, please take only Tylenol.

If you are particularly anxious about medical procedures, please discuss this with your provider. We may prescribe a mild sedative for use prior to the procedure, if necessary.

Eat a nutritious breakfast/lunch prior to the procedure. Maintain adequate hydration prior to appointment.

Shower before your visit. Suture-strips will be applied the day of the procedure and may be removed on day 4 following the procedure. You will be able to shower the morning following your procedure. Prior to the shower, remove the overlying gauze and you can shower with the underlying suture-strips in place.

Please avoid scraping/removing scab/shaving at the laser entry site(s).

Wear loose fitting, comfortable clothing. When you arrive home, we ask that you complete three separate 20 minutes walks during the day of the procedure and 1-2 walks a day for the following week. It is normal to notice a small amount of fluid drainage on the gauze following the procedure (in the initial 24 hours). Your leg may also feel "tight" and some bruising and discomfort is expected for 7-14 days afterwards.

Please bring your compression stockings to the office. We will only require you to wear stockings on the leg we treat. You will be asked to wear the compression stocking overnight, and then **during the day** for 5 days following the procedure.

Please arrange for someone to drive you home. This is especially important if you require a mild sedative with the procedure. You will be receiving local anesthetic during the laser procedure. Expect to be in office for approximately 1 hour.

You may return to work the next day. Some patients (in more intense occupations) find it helpful to take a day or two off of work. Please limit exercise to walking for 5-7 days following the laser. Running, weightlifting, and other high impact activities may be resumed at that time.

Please avoid flying for 5-7 days and long car trips (>2-3 hours) following the procedure.

Please avoid use of iron containing supplements during the first 4-6 weeks of your treatment protocol.

Please refrain from hot tubs/swimming for 5 days following the procedure.

Please have Motrin/Advil (ibuprofen 200mg-take three tablets every eight hours with food) **OR** Aleve (naproxen 225mg -take two tablets every 12 hours with food) available at home. You will be encouraged to use **ONE** of these medications on a daily basis for 1 to 2 weeks. Please let Dr. Schoonover or his team know if you are already taking a prescription **anti-inflammatory** like **Celebrex** or **Mobic**. Although unusual (<2% of patients), you may require an additional narcotic medication.

You will be scheduled for an appointment approximately 1 week after the procedure. A follow-up ultrasound will be performed at that time. Additional procedural appointments will also be scheduled as indicated with your individualized treatment plan.

Remember: Dr. Schoonover and his team may be reached after hours at 317.348.3020.

Better Options. Healthier Legs.®

www.indyveins.com

317.348.3020